



A Review on Aeromycological Studies

M.B. Kanade, Sujit Wagh, B.S. Mali and S.J. Chavan

P.G. Research Center, Department of Botany, Tuljaram Chaturchand College of Arts,
Science and Commerce, (Autonomous), Baramati Dist. Pune - 413 102, Maharashtra, India

*Corresponding author e-mail : mahadevkanade1@gmail.com

The present attempt has made to study the fungal flora available in the air, over agricultural fields and its impact on crop health. Furthermore this work also emphasizes on origin of the aeromycological research in India and abroad, past and current scenario of aeromycological work at national and International level, aeromycological studies over agricultural fields from India etc.

Keywords : Aerobiology, Aeromycology.

INTRODUCTION

Aerobiology studies small particles from biological origin in the air. Aeromycology is branch of Botany (Mycology) which deals with study of fungal flora in air. Aeromycology studies not only include fungal spores liberation from the sources transport and deposition, but also their effects on plants, animals, humans and even over food, building, work of art etc. Numbers of microscopic fungal spores are occurred in indoor and outdoor environment are commonly called as fungal bioaerosols. Singh and Dahiya (2008) reported more than 80000 fungal spores in air along with their dispersal mechanism and seasonal variation.

Fungal spores are dominating the other bioparticles in the air spora and fungal spores are mostly pathogenic, non-pathogenic as well as saprophytic in nature, living on dead and decaying organic remains of plants (Agrawal *et al.*, 1969). Fungi generally cause major diseases in plants. The importance of airborne micro fungi has long been recognized, especially in plant pathology. The disease incidence was greatly influenced by quantitative assessment of pathogens that is, fungi their prevailing meteorological parameters for disease development between the growth stages of the host plant and presence of pathogen.

Due to small size, fungal spores remain suspended in the atmosphere for a long time and causes number of plant diseases, respiratory disorders (Woolcock *et al.*, 2001) and many other allergic ailments. Kasprzyk (2008) stated that, knowledge of concentrations of airborne fungal spores is especially important for agricultural and occupational medicine. Aeromycology has its application in agrobiolgy, particularly with respect to pathogenic fungi and in the conservation of the artistic heritage.

The propagative phase of fungi is spores of which many are adapted to air transport. In many species, however, the spores rarely get into the air and are carried by insects or are either seed or soil borne. Fungal spores often constitute a major component of airspora. The spores are extremely variable in shape, size and colour with or without ornamentations on the spore wall. Most of these range from 3-20µm while a few are upto 100µm diameter (Tilak, 1982).

Pathogenic fungal bioaerosols are responsible to cause severe plant diseases to crop plants and they reduce crop yield (Karne, 2013). Stakman and Christensen (1946) made an intensive study and established the relationship between aerobiology and plant diseases and emphasized its role in forecasting the disease. The study of aeromycology is important in plant pathology and in disease forecasting of plant diseases (Lohare and Kareppa, 2010). The study of fungal aerospora is important to understand the dissemination of pathogenic spores in the atmosphere and successful disease management.

Origin of the aeromycological research : Air is the way fungi use to transports their spores. Many fungi are crop pathogens and cause important economic losses, hazards. Airborne fungi can be migrated thousands of kilometers away from their sources. One of the most common problems to humans is allergy to fungi spores. By the contrary of pollen grains fungi spores are frequent in indoor environment. About 100 species of fungi regularly involved in human and animal mycoses. Toxic secondary metabolite produced by fungi and their production depend on the environmental conditions (Ana *et al.*, 2002). The study of fungal air spora is important in order to understand its dissemination and spread in the atmosphere. Adequate information regarding the time and mode of the spread of the pathogens in air plays most important role in disease management of plants (Mishra and Srivastava, 1972). By keeping this in mind, it was thought to



study the effect of all these factors on the vegetation as well as on the microbes and their impact on health related problems of human being is utmost important.

Past and current scenario of aeromycological work:

International level : Meier *et al.* (1933) of U.S. and Stepanov (1935) of U.S.S.R. was established Aerobiology as a special branch. The honour of pioneering experimental aerobiology goes to Miquel (1879) who explain the technique to analyse the microbial air population. He produced the first suction trap and noted its improved collection efficiency. Roger and Meir (1936) reported the collection of microorganisms from 36,000 feet. Rittenberg (1939) investigated the microbiology of marine air.

According to Roger (2003) fungi affect humans in complex ways and are capable of eliciting a number of disease responses, such as infectious, allergi: and irritant and toxic effects. Fungal exposure is unequivocally associated with exacerbations of asthma, although the role of fungi in causing the disease is yet to be determined. The association between home dampness and respiratory health effects is strong and fungal exposure is suspected to be associated with the linkage.

Different microorganisms such as fungi, algae and bacteria may have a negative effect on the preservation of artistic-historical heritage, especially when microclimatic conditions favour their development. Numerous museum complexes therefore employ systematic temperature and humidity monitoring, with the aim to prevent or slow down their growth (De Nuntiis *et al.*, 2004).

Many researchers also report interaction between microscopic fungi and arthropods on the surface of wall paintings, which increases their alteration (Hoffland *et al.*, 2004). These biological agents affect not only the aesthetical appearance but also the structure of materials. Contributing to their preservation therefore is the main goal of studies of the mycobiota carried out in heritage buildings (Aira *et al.*, 2007).

Meier *et al.* (1933) reported spores in air of United States. Meier (1933) carried out collecting microorganisms from the Arctic atmosphere. Ainsworth (1952) investigated the incidence of airborne *Cladosporium* in the London region. Paddy *et al.* (1955) studied fungi in air over the Atlantic Ocean. Gregory and Hirst (1957) observed the summer air-spores at Rothamsted in 1952. Gregory (1973) reported the microbiology of the atmosphere of England during 1961-1973. Frey and Duris (1962) carried out the studies of air borne fungal spores by slide and culture methods. Lacey (1962) reported the summer air-spores of two contrasting adjacent rural site. Moustafa and Kamel (1976) investigated the fungal spore diversity from Kuwait and Levetin and Horowitz (1978) noticed from Tulsa. Larsen (1981) studied fungal aerospora from Copenhagen and Al-Doory *et al.* (1982) from Washington. Lyon *et al.* (1984), Abdel-Hafez and El-Said (1989), Nussbaum (1990), Larsen and Gravesen (1991) and Abdel- Hafez *et al.*

(1993) reported the effect of weather conditions on population of fungal aerospora.

Burge and Solomon (1978) reported sampling and analysis of biological aerosols. Banerjee *et al.* (1987) carried out indoor fungal flora from Durham, USA. Harriet *et al.* (1987) reported sampling and analysis of biological aerosols. Frinking (1991) carried out aerobiology of "closed" agriculture system. Shaheen (1991) reported aeromycology of Jordan. Rosas *et al.* (1992) carried out seasonal distribution of *Aspergillus* in the air of Mexico. Tan *et al.* (1992) observed variation in tropical airspora in Singapore. Sorlini (1993) has made aerobiology as general and applied aspects in the conservation of art works. The most frequent genera *Cladosporium*, *Alternaria*, *Aspergillus* and *Penicillium* in an urban and rural environment of East-France were noticed by Joel-Simeroy *et al.* (1993). Fungal airspora of Durban: a sub-tropical, coastal South African city were studied by Dames and Cadman (1994). Hirst (1994) carried out aerobiology at Rothamsted. Halwagy (1994) studied fungal airspora of Kuwait city during 1975-1987.

Past scenario of aeromycological work : National level Cunningham (1873) is the pioneer member of aerobiological research in India. He published his book entitled "Microscopic Examination of Air". Prof. K. C. Mehta extensively investigated the wheat rust spores from air. Furthermore, he observed that rust spores were incapable of surviving during high temperature of summer in the plains of North India. The present knowledge of rust disease of cereals is only due to his researches. Among the plant pathogen wheat rust is the best aerobiologically investigated one. Later on Rajan *et al.* (1952) noticed the fungal spora from Kanpur.

Padmanabhan *et al.* (1953) studied the incidence of conidia of *Helminthosporium oryzae* in paddy field at Cuttak. Kalra and Dumbrey (1957) studied airspora of Army MedicalCampus, Pune. Airspora over potato fields at upper Shilong has been studied by Konger and Baruah (1985). Gupta *et al.* (1960) reported airborne spores from Jaipur. Dua and Shivpuri (1962), Karnik (1962) recorded atmospheric pollen and fungal spores at Delhi and Jalgaon respectively. Sreeramulu and Seshavataram (1962) studied spore content of air over paddy fields at Pentapadu (A.P.). Nair (1963) investigated atmospheric pollen and fungal spores and other vegetable matter at Vellore, Madras. Chaubal and Deodikar (1964) studied airborne spores around Poona.

Prof. S. T. Tilak is the pioneer worker in aerobiology from Maharashtra. He discovered and developed many air samplers for quantitative study of air spora like Tilak Air Sampler, Volumetric Continuous Tilak Air Sampler and Rotorod Tilak Air Sampler. Furthermore he published many books on aerobiology viz. Aerobiology in 1982, Air Monitoring in 1987, Atlas of Air Borne Pollen Grains and Fungal Spores in 1989 and Aerobiology revised edition in 1995.

Anand *et al.* (1981) investigated the fungal spora from Bangalore city and he interrupted this information in



immunotherapy. Bhalke (1981) studied the aerobiological survey over some agricultural fields at Aurangabad. Dixit and Gupta (1981) noticed seasonal and diurnal fluctuation of airspora from barley field. Gaur and Kasana (1981) investigated atmospheric microbial population of Modinagar. Jankibai and Subba Reddy (1981) studied airspora of Vishakhapatnam. Tilak and Bhalke (1981) studied aeromycology at Aurangabad. Narayan *et al.* (1982) observed the atmospheric microbes of Malleswaram market at Bangalore. Agashe (1983) observed the airspora of Bangalore. Babu (1983), Bhosale (1983), Patil (1983), Saibaba (1983) and Wankhede (1983) recorded airspora at Aurangabad.

Mutha Reddy and Ramanujan (1989) studied the aerobiology of Hyderabad (A.P.). Bhat and Rajasab (1988) recorded the airborne fungal spores from Gulberga, Karnataka. Kavishwar (1990) recorded aerobiological studies at Dhulia. Kotwal (1991) observed the airspora over grapevine yard at Nasik. Pandey and Tiwari (1991) conducted an extensive aerobiological survey at Raipur (C.G.). Mishra and Zamil (1991) studied indoor environment fungi from mills of Lucknow. Verma and Chile (1992) studied the seasonality and allergenic behaviour of certain fungi at Jabalpur.

Aeromycological studies over agricultural fields from India

: The information on airborne allergenic fungal flora in rural agricultural areas is largely lacking. Adequate information is not available to the bioaerosol researchers regarding the choice of single versus multiple sampling stations for the monitoring of both viable and non-viable airborne fungi. There is no long-term study estimating the ratios of viable and non-viable fungi in the air and earlier studies did not focus on the fractions of airborne allergenic fungi with respect to the total airborne fungal load (Adhikari *et al.*, 2004).

Gadekar (2014) investigates the fungal aero-spora over *Sorghum* crop at Pune and reported that *Curvularia*, *Cladosporium*, *Helminthosporium*, *Cercospora*, smut, rust, *Nigrospora* and *Alternaria* were the most dominant mycoflora. Common occurrence of *Chaetomium* spores over jowar fields were investigated by Kavishwar (1990). Most common occurrence of *Alternaria* spores over jowar fields were studied by Navi *et al.* (2005) and Kanade *et al.* (2018a). Singh and Banopadhyay (2000) reported *Curvularia* spores were dominant over jowar fields. Sreenivasa *et al.* (2010) noticed common occurrence of *Aspergillus* along with *Sorghum* grains. Karne (2013a) reported 46 fungal spores over jowar fields and out of which 20 types (74.4%) were recorded as pathogenic and 25 types (81.5%) were recorded as allergenic. Patil and Mali (2017) investigated that *Alternaria*, *Plasmopara*, *Physarum*, uredospores, *Leptosphaeria* and *Cladosporium* spores were the most common over the *Sorghum* fields. Furthermore they reported that, the spore concentration of Deuteromycotina was found to be dominant. Aeromycoflora over the jowar fields from Barshi area were studied by Patil *et al.* (2016, 2016a) and Patil and Mali (2018) and reported 20 types of Ascomycotina fungi as dominant fungal flora during kharif season.

According to Karne (2013a) many plant pathogenic fungi viz. *Alternaria*, *Puccinia*, *Cercospora*, *Helminthosporium*, *Colletotrichum*, smut spores, *Diplodia*, *Sclerospora*, *Stemphylium* and *Fusarium* were recorded almost all round the year. Further reported that 50 fungal spores over wheat fields and out of which 21 types (71.7%) were recorded as pathogenic and 25 types (85.8%) were recorded as allergenic. Badge and Mhatre (2017) studied the airborne fungal spores over some paddy fields in the Ratnagiri district of Maharashtra and recorded that Deuteromycotina contributed highest percentage and lowest percentage contribution was found to be that of Zygomycotina to the total airspora. Nasrin Begum and Sudhendu Mandai (2017) studied the aeromycoflora diversity over a paddy field in Birbhum district of West Bengal and reported the seasonal variation of fungal spores occurred highest in winter season followed by rainy and summer season. Further noticed that *Alternaria* sp. showed the highest percentage of occurrence followed by *Curvularia* sp., *Helminthosporium* sp., *Cladosporium* sp., *Nigrospora* sp., *Aspergillus fumigatus* and *A. niger*.

Bagwan (2010) reported *Aspergillus* species were dominantly occurred from the vegetable market of Udgir. Kumar *et al.* (2013) noticed that, *Aspergillus* sp., *Penicillium* sp. and *Alternaria* sp. spores were most abundant in vegetable market environment of Hapur, Uttar Pradesh. Concentration of air borne fungal spores in main fruit and vegetable market of Agra were studied by Garg *et al.* (2015) and reported *Aspergillus flavus* was found to be most dominant and frequent mould in the aerospora. Frequent occurrence of *Alternaria* was studied by Vijayalaxmi *et al.* (2001) over chilly, Kumar and Kolte (2006) over mustard and Saha *et al.* (2006) over brinjal fields. Lohare and Kareppa (2010) noted that Deuteromycetes dominated all other groups and its mean contribution was 71.76% over onion fields at Udgir (Maharashtra). The most predominant species found in store houses of onion were *Aspergillus niger*, *A. flavus*, *Botrytis* sp., *Fusarium oxysporum*, *Helminthosporium* sp., *Trichoderma* sp., *Mucor mucedo* etc. and from ginger were *A. niger*, *F. oxysporum*, *F. solani*, *F. zingiberi*, *Lasiodiplodia* sp and *Verticillium thebromae* reported by Juri Devi *et al.* (2010). According to Kapadi and Patel (2019) weather conditions play an important role in seasonal development of many plant diseases in tomato fields (*Lycopersicon esculentum* Mill.). Further they reported that the high concentration of air born pathogenic fungal spores were responsible in diseases incidence in tomato crop e.g. early blight (*Alternaria solani*), late blight (*Phytophthora species*), *Fusarium* stem rot (*Fusarium species*).

According to Thakur (2016) the meteorological parameters during monsoon have found to play significant role in determining the spore load in the air over *Phaseolus vulgaris* L. fields from Pune. Furthermore recorded that Deuteromycotina spores have been highest during all the months as compared to other groups of fungi.

Aeromycological investigation over a pomegranate fields at Ahamdpur district were studied by Kadam *et al.*



(2010) and total 61 fungal spore types were recorded and *Ustilago*, *Cercospora*, *Helminthosporium*, *Colletotrichum*, *Fusarium* and *Alternaria* as pathogenic to pomegranate were noticed. Singh *et al.* (1992) reported fungal air spora over a pomegranate fields from Imphal and concluded that, the diseases appeared in the November, probably due to temperature 25-28°C. Highest concentration of Deuteromycetes spores (69.98%) to the total aerospora from pomegranate fields of Nashik were reported by Ahire *et al.* (2010). *Acaulopage*, *Albugo*, *Mucor* and *Rhizopus* noticed as chief fungal flora over pomegranate fields by Kadam and Khandia (2018). According to Aher *et al.* (2015) in the last several years fungal diseases has become a major concern to the pomegranate (*Punica granatum* L.) growers. Furthermore reported various air-borne fungal diseases of pomegranate like leaf spot caused by *Curvularia* and *Cercospora*, fruit rot caused by *Colletotrichum*, burning of fruits and leaves by *Alternaria alternate*. Deuteromycotina spores *viz.* *Alternaria*, *Aspergillus*, *Cercospora*, *Colletotrichum*, *Fusarium*, *Gloeosporium* and *Helminthosporium* were found dominantly and frequently over pomegranate fields (Kanade *et al.*, 2018a).

Babu (1983) investigated that *Hypoxyylon* spores were most common over banana fields at Aurangabad. In Nashik, *Cercospora* contributed 5.27% over the groundnut fields were observed by Sonawane (2013). Kshirsagar *et al.* (2016) reported 58 spores over groundnut fields of Seed district and noticed that Deuteromycotina was dominated the air spora. Aerobiological investigations were conducted for two consecutive kharif seasons of 2012 and 2013 at Newasa, Dist. Ahmednagar over ground nut fields by Sonawane *et al.* (2017) and reported that *Cladosporium* spore type was found to be the most dominant, followed by smut spores, *Alternaria*, *Aspergillus* and *Cercospora*.

An aeromycological survey over sunflower filed from Kada, Dist. Seed was under taken by Pathare *et al.* (2010). Furthermore reported that during the period of investigation the spores of *Alternaria* were almost continuously found in the atmosphere and this fungal spore contributed 13.75% width total concentration of 70364/m³ of air. Kshirsagar and Pande (2012) found the prevalence of spores of *Cladosporium* over sunflower fields from Rajuri. Allapure (2016) recorded 32 fungal spore types from sunflower field in Udgir region and noticed dominant spore type as *Cercospora* (13.22%), *Aspergillus* (9.52%), *Alternaria* (7.34%), *Nigrospora* (7.34%), *Leptosphaeria* (7.16%), rust spores (6.58%), *Cladosporium* (4.23%), *Claviceps* (4.23%), smut spores (4.23%), *Fusarium* (3.29), *Curvularia* (3.23%), *Bitrmonospora* (2.46%) and basidiospore (2.46%).

According to Kanade *et al.* (2018) *Alternaria*, *Mucor*, *Aspergillus*, *Cladosporium*, *Helminthosporium*, smut spores, conidia, chlamydospores or thalospores were observed as extremely abundant over sugarcane fields. Aeromycological investigation over a sugarcane fields at Ahamdpur district were studied by Kadam *et al.* (2010) and during the investigation total 61 fungal spores types were recorded and

Ustilago, *Cercospora*, *Helminthosporium*, *Colletotrichum*, *Fusarium* and *Alternaria* as pathogenic to sugarcane were recorded. Aeromycological survey of air borne fungi over sugarcane fields were done by Nanda and Nayak (2015) at Paducheri (Chennai) and recorded 18 fungal species belongs to 10 genera among the recorded fungi Deuteromycotina members were found as predominant followed by Zygomycotina and Ascomycotina. Singh *et al.* (1992) reported fungal air spora over a sugarcane fields from Imphal and concluded that, the diseases appeared in the November. It was probably due to temperature 25-28°C and occurrence of the smut spores was 16.38%. Occurrence of *Dendryphiopsis* spores over sugarcane fields were reported by Bhagwan (1983) from Nanded. Highest concentration of Deuteromycetes spores (69.98%) to the total aerospora from sugarcane fields of Nashik were reported by Ahire *et al.* (2010). *Acaulopage*, *Albugo*, *Mucor* and *Rhizopus* noticed as chief fungal flora over sugarcane fields by Kadam and Khandia (2018).

Cercospora sp., *Helminthosporium* sp., *Ramularia areola* and *Alternaria* sp. reported pathogenic to the cotton crop by Jaganath and Gaikwad (1998) at Ahmedpur, Maharashtra. Bembrekar and Bodke (2013) noticed 17 types of fungal spores belonging to the Ascomycetes over cotton fields during the kharif seasons from Nanded.

CONCLUSION

As fungal spores are microscopic they remain suspended in the atmosphere for a long time and causes number of plants, animals and allergic diseases (Woolcock *et al.*, 2001). Many researchers of aeromycology unanimously agreed with the chief contribution of Deuteromycotina spores in the atmosphere of majority all types of agricultural fields (Kanade *et al.*, 2018a). In India *Alternaria*, *Rhizopus*, *Helminthosporium*, *Aspergillus*, *Cladosporium* and *Cutvulerie* are allergic fungi and cause various diseases in plants as well as in humans (Shivpuri, 1982). Karne (2011) noticed that *Alternaria*, *Puccinia*, *Cercospora*, *Helminthosporium*, *Colletotrichum* and *Fusarium* were plant pathogenic fungi and found throughout the year over agricultural fields and infects the healthy crop. In Marathwada region percentage contribution of *Helminthosporium* was reported by Vaidya (1990) were 3.07 and by Jadhav (1990) were 2.23 from agricultural fields. Bisht *et al.* (2002) investigated that smut spores were most commonly observed over many cereal fields and grasses. Fungal diseases cause rotting of fruits, spot diseases of leaves and fruits and fruits discolouration etc. Ultimately there is reduction in the prize of the fruit in the market. *Aspergillus* was the most frequent genus along with *Cladosporium*, *Penicillium* and *Alternaria* spores were also fairly abundant from fruit market environment of Nagpur (Kakde *et al.*, 2001). Therefore, study of aeromycology is utmost important to understand the broadcasting of allergic and pathogenic spores and their successful disease management.



REFERENCES

- Abdel-Hafez, S. I., Abdel-Aal, H. M. and Ahmed, B. 1993.** Seasonal variations of fungi of outdoor air and sedimented dust at Assiut region, Upper Egypt, Grana, **32 (2)** : 115 - 121.
- Abdel-Hafez, Sobhyand El-Said, and Ahmed, B. 1989.** Seasonal Variations of Airborne Fungi in WadiQena, Eastern Desert, Egypt. Grana, **28** : 193 - 203.
- Adhikari, A., Sen, M. M., Gupta-Bhattacharya, S. and Chanda, S. 2004.** Airborne viable, non viable and allergenic fungi in a rural agricultural area of India: A two year survey at five outdoor sampling stations. Science of the Total Environment, **326** : 123 - 141.
- Agarwal, M.K., Shivpuri, D.N. and Mukherji, K.G. 1969.** Studies on allergenic fungal spores of Delhi, India metropolitan area (Botanical aspects). J. Allergy, **44** : 193 - 203.
- Agashe, S.N. 1983.** Aeromycoflora of Bangalore. Aspects of Allergy and Applied Immunology, 16.
- Aher, S.K., Dhawale, V.P. and Baviskar, P.S. 2015.** Qualitative assessment of airborne deuterospores over pomegranat (*Punica granatum* L.) field. Int. J. of Life Sciences, Special Issue **A3** : 18 - 20.
- Ahire, P. P., Kadam, V.B. and Patel, S.I. 2010.** Atmospheric concentration and seasonal variation in the smut spores at Nashik, Maharashtra, India. Plant Archives, **10 (2)** : 963 - 964.
- Ainsworth, G.C. 1952.** The incidence of airborne *Cladosporium* in London region. J. Gen. Microbio, **7** : 358 - 361.
- Aira, M.J., Jato, V., Stchigel, A.M., Rodriguez-Rajo, F.J. and Piontelli, E. 2007.** Aeromycological study in the cathedrals of Santiago de compostela (Spain). International Biodeterioration and Biodegradation, **60 (4)** : 231 - 237.
- AL-Doory, Y., Darson, J.F. and Best, J. 1982.** Further studies on the airborne fungi and Pollens of Washington, W. C. Metropolitan area. Ann. Allergy, **49** : 265 - 269.
- Allapure, R.B. 2016.** Aeromycological studies over sunflower field at Udgir region (M.S.). International Journal of Research in Medical and Basic Sciences, **02 (11)** : 49 - 52.
- Ana M. Calvo, Richard A. Wilson, Jin Woo Bok and Nancy P. Keller. 2002.** Relationship between secondary metabolism and fungal development. Microbiol. Mol. Biol. Rev. **66 (3)** : 447 - 459.
- Anand, P., Agashe, S. N. and Manjunath, K. 1981.** Aerobiological studies of Bangalore City and their scope in Immunotherapy. Proc. Nat. Conf. Biopollution, 117 - 124.
- Babu, M. 1983.** Aerobiological studies at Aurangabad, Ph.D. Thesis, Dr. B. A. Marathwada University, Aurangabad.
- Badge, M. and Mhatre, K. 2017.** Aeromycological studies over some paddy fields in the Ratnagiri district of Maharashtra State. International Journal of Current Research, **9 (06)** : 52441 - 52444.
- Bagwan, N. B. 2010.** Seasonal variation in Aeromycoflora of vegetable market at Udgir, Maharashtra, India. J. Mycol. Pl. Pathology, **40 (3)** : 360 - 364.
- Banerjee, U.C., Weber, P., Ruffin, J. and Banerjee, S. 1987.** Airborne fungi of some residences in Durham, North Carolina, USA. Grana, **26** : 103 - 108.
- Bembrekar, S.K. and Bodke, S.S. 2013.** Studies on composition and components of airspora belonging to ascomycetes over cotton field during two consecutive kharif seasons. Indian Streams Research Journal, **3 (2)** : 1 - 9.
- Bhalke, S.P. 1981.** Airspora over *Sorghum* fields. Ph.D. Thesis, Dr. B. A. Marathwada University, Aurangabad.
- Bhat, M.M. and Rajasab, A.H. 1988.** Incidence of viable fungal airspora in Gulberga during 1984-86, A two year survey. Indian Journal of Aerobiology, **1 (1)** : 10-16.
- Bhosale, S.S. 1983.** Studies in airspora of Aurangabad. Ph.D. Thesis, Dr. B. A. Marathwada University, Aurangabad.
- Bisht, V., Singh, B.P., Arora, N., Gaur, S.N. and Sridhara, S. 2002.** Antigenic and allergenic cross reactivity of *Epicoccum nigrum* with other fungi. Ann. Allergy Asthma Immunol., **89** : 285 - 291 .
- Burge, H.P. and Solomon, W.R. 1978.** Comparative merits of eight popular media in aerometric studies of fungi. Journal of Allergy and Clinical Immunology, **8 (3)** : 1199 - 203.
- Chaubal, P. D. and Deodikar G. B. 1964.** Air borne spores around Poona University. Journal of Poona University, **26** : 123 - 136.
- Cunningham, D. D. 1873.** Microscopic examination of air. Government Printer, Calcutta, 58.
- Dames, J. F. and Cadman, A. 1994.** Airspora of Darban, a Subtropical, Coastal South African city II fungal spore component. Grana, **33** : 346 - 348.
- De-Nuntis, P., Bitelli, L., Guaraldi, P., Monco, A. and Salvi, A. 2004.** In: Proceedings XI Internat. Palynological Congress (A. Polen, ed.), University of Cordoba, Cordoba, 88.
- Dixit, R.B. and Gupta, J.S. 1981.** Seasonal and diurnal concentration of aerospores over barley field. Journal of Indian Botanical Society, **60 (4)** : 348 - 351.
- Dua, K. L. and Shivpuri, D. N. 1962.** Atmospheric fungal spores in delhi in 1958-1959. J. allergy, **3** : 507 - 512.
- Frey, D. and Durish, S. 1962.** Estimation of air-borne fungal spores- A Comparison of slides and culture methods. Mycopath Mycological Applications. **16** : 295 - 303.
- Frinking, H. D. 1991.** Aerobiology of "closed" agricultural systems. Grana, **30 (2)** : 481 - 485.
- Gadekar, S.S. 2014.** Diversity of fungal spores over jowar crop. Int. J. of Life Sciences, **2 (2)** : 155-159
- Garg, A., Singh, R. and Singh, S. 2015.** Incidence of toxigenic isolates of *Aspergillus flavus* in the aerospora of main fruit and vegetable market at Agra. Nature and Environment, **20 (1)** : 33 - 36.
- Gaur, R.D. and Kasana, M.S. 1981.** Studies on aerobiology of Modinagar. Journal of Indian Botanical Society, **60** : 266 - 277.
- Gregory, P. H. and Hirst, J. M. 1952.** The Summer Air-Spora at Rothamsted. J. Gen. Microbiol. **17** : 185 - 152.
- Gregory, P. H. 1973.** The microbiology of the atmosphere New york 2nd ed., 337.
- Gupta, K. D., Sogani, J. L. and Kasiwal, R.M. 1960.** Survey of the allergenic aerial mold spores at Jaipur. Indian J. Gen. Microbiol. **20** : 328 - 354.
- Halwagy M. H. 1994.** Fungal Airspora of Kuwait City, Kuwait, 1975-1987. Grana, **33 (6)** : 340 - 345.
- Harriet, A. Burger and William R Solomon, 1987.** Sampling and analysis of biological aerosols. Atmospheric Environment, **21 (2)** : 451 456.
- Hirst, J. M. 1994.** Aerobiology at rothamsted. Grana, **33 (2)** : 66 - 70.
- Hoffland, E., Kuyper, T. W., Wallander, H., Plassara, C., Gorbushina, A. A., Haselwandter, K., Holmstrom, S., Landeweert, R., Lundstrom, U.S., Rosling, A., Sen, R., Smits, M. M., Van Hees, P. A. W. and Van Breemen, N. 2004.** The role of fungi in weathering. Frontiers in Ecology and the Environment, **2** : 258 - 264.
- Jadhav, D. S. 1990.** Aerobiology of groundnut at Kallam. Ph.D. Thesis, Dr. B. A. Marathwada University, Aurangabad.
- Jagannath, S. and Gaikwad, Y. B. 1998.** Aerobiological approach to aerolate mildew disease of cotton. Ind. J. Aerobiol., **11 (1 & 2)** : 33 - 37.
- Janakibai, A. and Subba-Reddy. 1981.** The Fungal Airospora of Visakhapatnam. Proc. Indian Natural Science Academy. **47(5)** : 731-747.
- Joel, S., Jean-Pierre, C. and Danielle, L. 1993.** Seasonal variations in the airborne fungal spore population of the East of France. Comparison between urban and rural environment during two years. Aerobiologia, **9** : 201 - 206.



- Juri Devi, Sadhana, Medhi and Sarma, T. C. 2010.** Aeromycological study of store houses of onion and ginger in Guwahati. The Bioscan, **2** : 547 - 552.
- Kadam, S.S. and Khandia, R. 2018.** Studies on composition and components of airspora belonging to phycocomycetes over the field of sugarcane. International Journal of Recent Trends in Science and Technology, Special Issue, ICRAFHN 2018,128-130.
- Kadam, R.M., Reddy, N.J.M. and Biradar, R.P. 2010.** Aeromycoflora over a sugarcane field at Ahmedpur Dist. Latur (M.S.). International Journal of Plant Sciences, **5 (2)** : 587 - 589.
- Kakde, U. B., Kakde, H. U. and Saoji, A. A. 2001.** Seasonal variation of fungal propagules in a fruit market environment, Nagpur (India). Aerobiologia, **17** : 177 - 182.
- Kalra, S. L. and Dumbrey, D. G. 1957.** Aerobiology of Army medical campus Poona (India). Armed Forces Med. J., **13** : 3 - 16.
- Kanade, M.B., Awatade, A., Gulave, A., Wagh, S., Kalkute, A., Chandankar, S. and Waghmare, A. 2018.** Aeromycological investigations over wheat, sugarcane and grape fields at Baramati (Pune), Maharashtra, Bioscience Discovery, **9 (1)** : 86 - 89.
- Kanade, M.B., Awatade, A., Gulave, A., Wagh, S., Kalkute, A., Chandankar, S. and Waghmare, A. 2018a.** Aeromycoflora over jowar and pomegranate fields at Baramati, Dist. Pune (M.S.), Bioscience Discovery, **9 (1)** : 93 - 96.
- Kapadi, M.R. and Patel, S.I. 2019.** Aeromycological approach of some fungal diseases on tomato crop (*Lycopersicon esculentum* Mill.) at Nashik, India. Journal of Drug Delivery and Therapeutics, **9 (3)** : 329 - 331.
- Karne, A.V. 2013.** Exploration of fungal biopollutants in the ambient air over groundnut field. Advances in Plant Sciences, **26 (2)** : 283 - 286.
- Karne, A. V. 2013a.** Aeromycological investigations in the ambient air over some crop fields in context to pathogenic and allergenic fungal bioaerosols. Nature Environment and Pollution Technology, **12 (4)** : 695 - 698.
- Karne, A.V. 2011.** An investigation of fungal aerobio-pollutants in the ambient air over maize fields. Nat. Env. Poll. Tech., **10 (4)** : 609 -612.
- Karnik, C. R. 1962.** A contribution to the rain water forms and air-spores of Jalgaon dist. Sci. and Cult., **28** : 475 - 476.
- Kasprzyk, I. 2008.** Aeromycology main research fields of interest during the last 25 years. Ann. Agric. Environ. Med., **15 (1)** : 1 - 7.
- Kavishwar, A. D. 1990.** Aerobiological studies at Dhulia-I (Jowar crop). Ph.D. Thesis, Dr. B. A. Marathwada University, Aijrangabad.
- Konger, G. and Baruah, H. K. 1985.** The incidence of air borne spores in potato plantation of supper shilling. J. Gauhati Uni., **9** : 81 - 89.
- Kotwal, S. G. 1991.** Aerobiological investigation at Nashik (airspora over Grapevine yard and Airspora inside Cattle shed). Ph.D. thesis, Dr Babasaheb Ambedkar Marathwada University, Aurangabad.
- Kshirsagar, J.J. and Pande, B.N. 2012.** Prevalence of Cladosporium spores over sunflower fields at Rajuri (N) M. S., India. Science Research Reporter, **2 (1)** : 66 - 68.
- Kshirsagar, J. J., Dharasurkar, A.N. and Pachkore G.L. 2016.** Fungal spore of groundnut fields in Beed District. International Journal of Researches in Biosciences, Agriculture and Technology, **4 (1)** : 161 - 164.
- Kumar, B. and Kolte, S. J. 2006.** Development of *Alternaria* blight in genotypes of Indian mustard (*Brassica juncea* (L.) Czern and Cross) under field. Indian Phytopathology, **39 (3)** : 314 - 317.
- Kumar, S., Sharma, S., Kumar, M., Sharma, P. K. and Sharma, N. 2013.** Seasonal variation of fungal propagules in vegetable market environment' of Hapur Uttar Pradesh India. International Journal of Microbial Resource Technology, **2 (1)** : 1- 6.
- Lacey, M. E. 1962.** The summer airspora of two contrasting adjacent rural site. J. Gen. Microbia., **29** : 485 - 501.
- Larsen, L. and Gravesen, S. 1991.** Seasonal variation of outdoor airborne viable microfungi in Copenhagen, Denmark. Grana, **30** : 467-471.
- Larsen, L.S. 1981.** A three year survey of microfungi in the air of Copenhagen. Allergy, **36** : 15 - 22.
- Levetin, E. and Horowitz, L. 1978.** A one-year survey of the airborne molds of Tulsa, Oklahoma. I. Outdoor survey. Ann. Allergy, **41** : 21-24.
- Lohare, S.D. and Kareppa, B.M. 2010.** Air spora over onion field. International Research Journal, **1 (3&4)** : 116 - 117.
- Lyon, F.L., Framer, C.L. and Eversmeyer, M.G. 1984.** Variation of airspora in the atmosphere due to weather conditions. Grana, **23** : 177-181.
- Meier, F.C., Stevenson, J.A. and Charles, V.K. 1933.** Spores in upper air. Phytopath., **23** : 23.
- Miquel, P. 1879.** Annual Report in Annuaire de l' observatoire de Montsouris.
- Mishra, R.R. and Srivastava, V. B. 1972.** Aeromycology of Gorakhpur - V. Airspora over wheat and barley fields, Mycopath et Mycol. Appl., **47** : 349 - 355.
- Moustafa, A.F. and Kamal, S.M. 1976.** A study of fungal spore population in the atmosphere of Kuwait. Mycopath, **59** : 29 - 36.
- Mutha Reedy, K. and Ramanujam, C.G. 1989.** An aerobiological study of Hyderabad (A.P.), Asian J. PL Sci. **1** : 7 - 21.
- Nair, P. K. 1963.** An analysis of atmospheric pollen, fungal spores and other vegetable matter of Vellore, Madras state. International J. Med. Res. **51** : 447 - 452.
- Nanda, A. and Nayak, B.K. 2015.** Studies on environmental bio-pollution by airborne fungi over a Sugarcane field. Der Pharmacia Lettre, **7 (5)** : 245 - 249.
- Narayan, M.C., Ravichandran, V. and Sullia S.B. 1982.** Aeromycology of the atmosphere of Malleswaram Market, Bangalore. Act. Bot. Indica, **10** : 196 - 200.
- Nasrin Begum and Sudhendu Mandai. 2017.** Aeromycoflora diversity over a paddy field in Birbhum district, West Bengal of Eastern India. J. Mycopathol. Res. **55 (2)** : 163 - 167.
- Navi, S. S., Bandyopadhyay, R, Reddy, R.K., Thakur, R.P. and Yang, X.B. 2005.** Effect of wetness duration grain development stages on *Sorghum* grain mold infection. Plant Diseases, **89** : 872 - 878.
- Nussbaum, F. 1990.** Variation in the Airborne Fungal Spore Population of the Tuscarawas Valley with Respect to Microenvironment, Time of Day, and Date, OHIO Journal of Science, **90 (3)** : 77 - 86.
- Paddy, S.M., Cramer, C.L. and Willey, B.J. 1955.** Fungi in air over Atlantic Ocean. Mycologia, **47** : 34 - 50.
- Padmanabhan, S.Y., Ganguli, D. and Balkrishnan, M.S. 1953.** *Helminthosporium* disease of Rice II. Source and development of seedling infection. Indian Phytopathology, **5** : 96 - 105.
- Pandey, P. and Tiwari, K.L. 1991.** Aeromycoflora of Raipur. (Abst.) 6 th Nat. Con. Aerobiol. Pandichery 113.
- Pathare, G.M., Mali, V.P. and Pande, B.N. 2010.** Atmospheric concentration of *Alternaria* spores over sunflower field. Inter. J. Plant Sci., **5 (1)** : 395 - 396.
- Patil, M.T. and Mali, N.S. 2018.** Seasonal variation in ascomycotina fungi over *Sorghum* crop at Barshi Tehsil of Solapur district Maharashtra. Bioscience Discovery, **9 (1)** : 76 - 78.
- Patil, C. R. 1983.** Aerobiological studies at Aurangabad. Ph.D. Thesis, Dr. B. A. Marathwada University, Aurangabad.
- Patil, M.T. and Mali, N.S. 2017.** Fungal Aerospora over the Jowar Fields. Central European Journal of Experimental Biology, **5 (1)** : 72 - 76.
- Patil, M.T., Kanade, M.B. and Mali, N.S. 2016.** Present Status of fungal diseases of jowar fields at Barshi area, Maharashtra. Proceedings of International Conference Plant Research and Resource Management organized by Tuljaram Chaturchand College, Baramati, Dist. Pune (M.S.), India, 166 - 168.



This document was created with Win2PDF available at <http://www.win2pdf.com>.
The unregistered version of Win2PDF is for evaluation or non-commercial use only.
This page will not be added after purchasing Win2PDF.




Principal
Tuljaram Chaturchand College
Baramati