



## SCREENING AND ISOLATION OF IMIDACLOPRID DEGRADING MICROORGANISMS FROM PESTICIDES CONTAMINATED SOIL

Vishakha N. Ganvir<sup>1</sup>, Ashok E. Kalange<sup>2</sup> and Shivaji J. Sathe<sup>3</sup>

<sup>1</sup>Department of Microbiology, Waghire College Saswad.

<sup>2</sup>Department of Physics, T.C.College, Baramati.

<sup>3</sup>Department of Microbiology, T.C.College, Baramati.

### ABSTRACT:

Pesticides are the chemical substances that prevent, kill and repel any pest. Repeated use of same pesticide, bulky handling or accidental release results in accumulation of pesticides residues or its metabolites in soil or water streams. The persistent nature of pesticides carries potential hazards to man and show lethal effects on living system. Imidacloprid (1-[(6-chloro-3-pyridinyl)-methyl]-N-nitro-2-imidazolidinimine), is a second generation neonicotinoid pesticide (chloronicotinyl insecticide). It is used to treat a wide range of pests on rice, maize, potatoes and vegetables. Imidacloprid has been reported as a stable compound in the environment with a half life over 100 days. There are scanty reports on imidacloprid degrading soil microorganisms. In an effort to find active microbial strains over the problem of commonly used pesticides like imidacloprid, a study was carried out. The objectives of this study were to screen, isolate and characterize the efficient imidacloprid degrading microorganisms from the contaminated agricultural soil. Imidacloprid degrading strains were enriched and isolated from vegetable farms of Harni and Jogawadi, Maharashtra. The minimal salt medium (MSM) was used for testing the imidacloprid degrading ability of the isolates. Among 20 soil isolates, four isolates showing maximum imidacloprid degrading ability were characterized on their morphological, cultural and biochemical characteristics. The characterized isolates were *Bacillus* spp., *Azotobacter* spp, *Actinomycetes* spp. and *Pseudomonas* spp. Imidacloprid was degraded by *Actinomycetes* spp up to 500 mgL<sup>-1</sup> and *Bacillus* spp., *Pseudomonas* species and *Azotobacter* species were upto 200 mgL<sup>-1</sup>. Detection of presence of NO<sub>2</sub><sup>-</sup> in the inoculated broth was the indication of imidacloprid degradation by the microorganisms. These findings suggest that these strains may be the promising organisms for bioremediation of imidacloprid contaminated soils.



**KEYWORDS:** Biodegradation, bioremediation, Imidacloprid and insecticide.

### 1.INTRODUCTION:-

The use of pesticide has become important in agriculture as pests destroy 30% of the agricultural product. India is the largest consumer of pesticide [1]. Pesticides are large and varied group of substances that

are specifically designed to kill biological organisms including weeds, insects and rodents. Pesticides include insecticides, fungicides, herbicides, rodenticide, molluscicides, nematocidal and plant growth regulators [2]. Classification of

pesticides can be done on the basis of chemical structure, physical state and target organism. The indiscriminate use of pesticides leads to the potential hazards to man and show lethal effects on living system [3]. Chronic pesticide

poisoning is attributed to increasing incidences of cancer, chronic kidney, suppression of immune system, sterility among males and females, endocrine disorders, neurological & behavioral disorders especially among childrens [4]. Insecticides like organochlorine and thyphus have been banned or restricted after 1960s in most of advanced countries [5].

Imidacloprid (1-[(6-chloro-3-pyridinyl)-methyl]-N-nitro-2-imidazolidinimine) is a second generation neonicotinoid insecticide (chloronicotinyl insectide). Imidacloprid is commonly applied as a systemic pesticide and effective against many sucking insects, fleas, termites and other chewing pests [6]. Because of its excellent systemic properties it can be used as seed dressing, as soil and foliar treatment in different crops including rice, maize, potatoes, cotton, cereals, sugarbeets and vegetables. In industry imidacloprid is considered to be a pesticide of relatively low toxicity, it has been found to be very toxic to the non-target insects such as bees [7, 8]. Sorption of imidacloprid increases with increasing soil organic carbon and pesticide concentration. The dissipation rates of imidacloprid in field are varied widely and it has been reported as a stable compound in the environment with a half life over 100days. In non-vegetated soil it has been reported with half life exceeding 180 days and more than three years in dry and aerobic conditions [9]. The degradation of imidacloprid in soil is generally facilitated by chemical, sunlight and microbial agents. Biodegradation is the most commonly used method [10]. Biodegradation studies of imidacloprid in soil demonstrated that imidacloprid urea, imidacloprid guanidine and 6-chloronicotinic acid are the possible metabolic products [11, 12]. Previous reports have revealed that very few species of bacteria are able to degrade this compound. The imidacloprid degradation by microorganism *Leifsonia* strain PC-21 is the first report [13].

The aim of the present work was to screen, isolate and characterize imidacloprid-degrading microbial strains which have the maximum ability to degrade high concentrations of imidacloprid.

## 2. MATERIALS AND METHODS

### 2.1. Chemicals

The technical grade imidacloprid was made available from the Chemistry department, Tuljaram Chaturchand College, Baramati.

### 2.2. Culture media

#### Mineral salt medium (MSM)

MSM was used for the isolation of bacteria [14]. The composition of the media in grams per litre of distilled water was Na<sub>2</sub>HPO<sub>4</sub> (2.4), KH<sub>2</sub>PO<sub>4</sub> (2), NH<sub>4</sub>NO<sub>3</sub> (0.1), MgSO<sub>4</sub>.7H<sub>2</sub>O (0.01), CaCl<sub>2</sub> (0.01) at pH 6.5. The medium was sterilized by autoclaving at 121 °C for 20 min.

### 2.3 Collection of soil sample

For the isolation of pesticide degrading microorganisms, soil sample was collected from Harni and Jogawadi, Maharashtra which was having history of repeated pesticide application. The soil sample collected in a sterile zipper polythene bags from 10cm of depth was air dried, sieved and stored at 4°C before processing for characterization.

### 2.4 Physico-chemical parameters of soil sample

Physico-chemical parameters were studied according to standard procedure [1].

**2.4.1. Moisture content:** 10gm of collected soil sample was dried at 60°C for 72h in oven. The dry weight of sample was taken till it showed constant weight. The % moisture was expressed as follows:

$$\text{Moisture \%} = \frac{W_1 - W_2}{W_1} \times 100$$

Where, W<sub>1</sub> = Weight of soil before oven drying

W<sub>2</sub> = Weight of soil after oven drying

**2.4.2. pH of soil sample:** Soil sample were dried at 60 °C for 72 h, powdered in pestle and mortar and filtered through 2 mm sieve and the sieved soil were dissolved in distilled water (2.5w/v) and vortexed for 5 minutes at 120 rpm then pH was measured by digital pH meter.

**2.4.3. Percent organic Carbon/Nitrogen:** One gram soil sample was mixed with 10 ml potassium dichromate (1 N) and 20 ml concentrated H<sub>2</sub>SO<sub>4</sub>. Then 150 ml distilled water and 25 ml FeSO<sub>4</sub> (0.5 M) were added and the excess was titrated against potassium permanganate (0.1N) solution to pink end point.

$$\% \text{ Organic carbon} = A - B \times 0.3 \times 1.33 / C$$

#### Where

A. Volume of K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> X Normality of K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub>,

B. Volume of KMnO<sub>4</sub> X Normality of KMnO<sub>4</sub>,

C. Weight of sample

Soil organic nitrogen was calculated using following equation:

$$\mathbf{2.4.4. \text{ Organic nitrogen (\%)} = 0.862 \times \% \text{ organic carbon}}$$

#### 2.5 Isolation and characterization of imidacloprid degrading bacteria: Primary screening:

5gm of collected soil sample was inoculated in 100ml of Mineral Salt Broth (MSB) containing 50mgL<sup>-1</sup> imidacloprid for bacterial isolation. Flasks were incubated at 37°C for 5-7days. Loopful of culture was streaked from enriched MS broth on sterile MSM plates containing pesticide and incubated at 37°C for 24-48h. Individual colonies were sub cultured on MSM containing the same concentration of pesticide. Bacterial colonies were characterized on the basis of morphology, cultural and biochemical tests like capsule staining (Manvel's method), Gram staining, motility, catalase, oxidase and sugar fermentation tests, IMViC test.

#### 2.6 Degradation of pesticide at different concentration by bacterial isolates

Soil bacterial isolates were streaked on MSM agar plates respectively at different concentrations of imidacloprid (50 mg L<sup>-1</sup> to 500mg L<sup>-1</sup>). Plates were incubated for 24-48h and growth was observed.

#### 2.7 Determination of imidacloprid metabolites by Nitrite detection test

Microorganisms were grown in MSM broth containing imidacloprid for 4-5days. Uninoculated MSM broth with pesticide and inoculated MSM broth without pesticide were taken as control. After incubation, 3-4 drops of two reagents sulfanilic acid and alpha Naphthylamine were added to 1ml of each inoculated and uninoculated broth and mixed.

### 3. RESULT AND DISCUSSION

**3.1 Physicochemical parameters of soil sample:** The physicochemical parameters affect the growth and diversity of microorganisms. The moisture content, certain amount of organic matter like carbon and nitrogen is essential for the growth and active functioning of the microbial population in the soil. Therefore it is necessary to study the soil factors and its relation to the microbial population. In the present study some of the important physicochemical properties of the soil were determined for the evaluation of natural microbial flora of the soil. ( Table1)

The analysis of physicochemical parameters of soil was done in the present study for the isolation of microorganisms. The colour of the soil sample was black with pH (7.0) and percent moisture content (0.62). The percent organic carbon and percent organic nitrogen was (0.3574) and (0.3167) respectively. These results were parallel to the findings of Rohilla SK et. al; (2012).

**3.2 Isolation of imidacloprid degrading microorganisms:** Primary screening was done by using enrichment technique. With successive transfers in MSM agar medium with imidacloprid as the only source of carbon and energy, bacterial strains were obtained from the soil sample degrading this

pesticide. On the basis of morphology, staining technique, biochemical characters and by referring Bergey's manual of determinative bacteriology, the isolated strains on MSM media were *Azotobacter spp*, *Actinomycetes spp*, *Bacillus spp* & *Pseudomonas spp*.

**3.3 Microbial degradation of imidacloprid at different concentration:** When these soil isolates were grown at different concentration of imidacloprid (50 mgL<sup>-1</sup> to 1000 mgL<sup>-1</sup>), *Actinomycetes spp*. were able to degrade imidacloprid up to maximum concentration of 500 mgL<sup>-1</sup> whereas *Bacillus*, *Pseudomonas species* and *Azotobacter species* were up to 200 mgL<sup>-1</sup> (Table 2). The result shown by bacterial isolates in this study were able to degrade imidacloprid at higher concentration than the findings of Geeta Negi et.al, 2014 where they found the tolerance of imidacloprid by bacterial isolates from 100 to 200 mg L<sup>-1</sup>.

**3.4 Determination of imidacloprid metabolites (imidacloprid guanidine) by Nitrite (NO<sub>2</sub>) detection test:** *Actinomycetes Spp*, *Azotobacter*, *Pseudomonas spp.* & *Bacillus spp.* have shown red colour after addition of 3-4 drops of both the reagents namely Sulfanilic acid and alpha Naphthylamine. Red colour was not detected in any of the control. Thus the formation of red colour indicates that nitrite is present in the broth inoculated with imidacloprid and microorganism. This indicates that these microbial species may be responsible for cleaving off the nitrite group on the end of the imidazolidin ring of imidacloprid to produce its metabolite imidacloprid guanidine. Research work of Jennifer C. et. al, (2007) have found concomitant production of nitrite as a metabolite of imidacloprid degradation by analytical procedure (cadmium reduction of nitrate to nitrite with colorimetric detection).

#### 4. CONCLUSION

Microorganisms isolated from contaminated soil were able to grow on media containing imidacloprid & degrade it by using imidacloprid as a sole source of carbon and energy. The highest pesticide degradation was found by *Actinomycetes spp* up to 500 mgL<sup>-1</sup> and *Bacillus spp.*, *Pseudomonas species* and *Azotobacter species* were upto 200 mgL<sup>-1</sup>. Presence of NO<sub>2</sub> was the indication of the imidacloprid degradation into its metabolite imidacloprid guanidine. The application of these microorganisms or microbial consortium can improve the bioremediation over conventional technique such as land rolling or incineration. Thus these microorganisms can be used to remediate soil contaminated with pesticides.

#### REFERENCES

1. Rohilla S.\*and Salar R. (2012). Isolation and Characterization of Various Fungal Strains from Agricultural Soil Contaminated with Pesticides. *Res.j.Recent.sci.1*:297-303.
2. Kavita Rani and Dhaniya G. (2014). Bioremediation and Biodegradation of Pesticides from Contaminated Soil and Water- a Novel Approach. *Int.j.Curr.Microbiol.app.Sci* 3(10):23-33.
3. Baishya K, Sharma H. (2014). Isolation and characterization of organophosphorus pesticide degrading bacterial isolates. *Arch. appl. Sci. Res.*6(5):144-149.
4. Agnihotri N. (1999). Pesticide: Safety Evaluation and Monitoring. All India coordinated Project (AICRP) on Pesticide Residues. Indian Agricultural Research Institute, New Delhi, India, 132-142.
5. Aktar M., Sengupta D, Chowdhary A. (2009). Impact of pesticides use in agriculture: their benefits and hazards. *Interdisc Toxicol* 2(1):1-12.
6. Nauen, Ralf, Denholm, Ian (2005). Resistance of insect pests to neonicotinoid insecticides: current status and future prospects. *Arch. Insect Biochem. Physiol.* 58, 200-215.
7. Sheets L. (2001). Imidacloprid: a neonicotinoid insecticide. In Krieger, Robert (Ed.), second ed., Handbook of Pesticide Toxicology, 2, 1123-1130 (Chapter 54).
8. Sladan S. Miroslav K., Ivan S., Snezana J., Petar K., Goran T., Jevdovic R. (2012). Resistance of Colorado potato beetle ( Coleoptera Chrysomelidae) to neonicotinoids, pyrethroids and nereistoxins in Serbia. *Rom. Biotechnol. Lett.* 17 (5), 7599-7609.

**9. Scholz K., Spiteller M. (1992).** Influence of groundcover on the degradation of <sup>14</sup>C-imidacloprid in soil. Brighton Crop Protection Conference- Pests and Diseases, 883-888.

**10. Bassey, D. E., Grigson, S.J.W. (2011).** Degradation of benzyldimethyl hexadecylammonium chloride by *Bacillus niabensis* and *Thalassospira* sp. Isolated from marine sediments. *Toxicol. Environ. Chem.* 93 (1), 44-56.

**11. Jesche P., Nauen R. (2005).** In Gilbert L., Iatrou K., Gill S.S. (Eds.), *Neonicotinic Insecticides Incomprehensive Molecular Insect science*, Elsevier.

**12. Krohn J., Hellpointer E. (2002).** Environmental fate of imidacloprid. *Pflanzenschutz-Nachrichten Bayer* 55, 1-26.

**13. Anhalt J., Moorman T., Koskinen W. (2007).** Biodegradation of imidacloprid by an isolated soil microorganism. *J. Environ. Sci. Health, Part B* 42, 509-514.

**14. Radehaus P., Schmidt S. (1992).** Characterization of a novel *Pseudomonas* sp. that mineralizes high concentrations of pentachlorofenol. *Applied and Environmental Microbiology* 58, 2879-2885.

**15. Negi G. Pankaj, Srivastava A., Sharma A. (2014).** In situ biodegradation of endosulfan, imidacloprid and Carbendazim using indigenous bacterial cultures of agriculture fields of Uttarakhand, India. *Int. J. of Biol, Biomol, Agri, Food and Biotechnol Engg.* 8 (9).

**Table 1. Physicochemical Parameters of soil sample**

Sample	Soil characteristics					
	Pesticide application	Colour	% Moisture	pH	Organic carbon %	Organic nitrogen %
Soil	History of 2yrs	Black	0.62	7.0	0.3574	0.3167

**Table 2. Degradation of imidacloprid at different concentration by cultures:-**

Organisms	Concentration in mg L <sup>-1</sup>						
	50	100	125	200	250	500	1000
<i>Actinomycetes spp.</i>	+	+	+	+	+	+	-
<i>Azotobacter spp.</i>	+	+	+	+	-	-	-
<i>Pseudomonas spp.</i>	+	+	+	+	-	-	-
<i>Bacillus spp.</i>	+	+	+	+	-	-	-

+ = Growth      - = No growth