



## SYNTHESIS AND CHARACTERIZATION OF SOME NEW THIAZOLYL SCHIFF BASE-METAL COMPLEXES

**Vaibhav P. Landage**

Department of Chemistry, Anekant Education Society's Tuljaram Chaturchand College, Savitribai Phule Pune University, Baramati, Pune, India

**Rahul S. Bhondwe**

Department of Chemistry, Anekant Education Society's Tuljaram Chaturchand College, Savitribai Phule Pune University, Baramati, Pune, India

**Bhausahab K. Karale**

Department of Chemistry, Rayat Shikshan Sanstha's Radhabai Kale Mahila Mahavidyalaya, Savitribai Phule Pune University, Ahmednagar, India

**Dilip R. Thube**

Department of Chemistry, Ahmednagar Jilha Maratha Vidya Prasarak Samaj's New Arts, Commerce and Science College, Savitribai Phule Pune University, Parner, Ahmednagar, India

Email: [drthube@rediffmail.com](mailto:drthube@rediffmail.com)

### ABSTRACT:

Our present work focused on the synthesis and characterization of Copper and Nickel metal complexes containing thiazolyl Schiff bases obtained by condensation of the ligands 2-(4-chlorophenyl)-4-methylthiazole-5-carbohydrazide and aromatic aldehyde by refluxing together in alcohol to get new series of thiazolyl Schiff base. The synthesized metal complexes were characterized by various spectroscopic techniques including elemental analysis, UV, FT-IR, <sup>1</sup>HNMR and XRD. Geometries of the Cu (II) and Ni (II) complexes were determined by indicated spectroscopic values. X-ray powder diffraction confirms that the complexes are crystalline in nature.

**KEY WORDS:** Schiff base, Thiazole, XRD.

### Article History

\* Received: 24/08/2021; Accepted: 16/09/2021

Corresponding author: Dilip R. Thube

### INTRODUCTION:

Schiff's bases and their derivatives have excellent physiochemical and biological properties. They are proven to be a promising industrial material and used in chemical sensors, optics and catalysis (Sahin et al., 2018), (Anacona et al., 2021; Bursal et al., 2021; Chaviara et al., 2004). They are an excellent drug candidate as antibacterial agents, fungicidal, antitumor and anticancer agents (Anacona et al., 2021; Damercheli et al., 2019; Manimohan et al., 2020; Shi et al., 2020; Singh & Barman, 2021). Diverse derivatization of transition metal Schiff's base complexes can be achieved due to the ease of synthesis by condensation of amines and aldehydes. These ligands coordinate with numerous metals and stabilize in several oxidation states. Schiff base ligands with heterocyclic moiety are "privileged ligands" due to their excellent therapeutic utility (Chaviara et al., 2004; Karatepe & Karatas, 2006; Pontiki et al., 2008; Yernale & Bennikallu Hire Mathada, 2014). Especially the ligands with N and S have drawn attention of many researchers due to the excellent chelation properties.

Thiazoles are an important class of heterocyclic compounds contains N and S atoms. Thiazoles are found in many natural products such as Dolastatin 10, Tubulysin, Largazole, Apratoxin [4] and in pharmacophore of many blockbuster drugs for instance sulphathiazole Fentinol (anti-inflammatory), Cambendazole (fungicide), Niridazole (schistozomicidal) and Ritonavir (anti-HIV, Covid-19) (Alsharif & Alam, 2017). Since last few decades researchers are attracted towards field of bioinorganic chemistry and coordination chemistry of thiazolyl Schiff's base metal complexes. Recently Cozzi nicely reviewed the importance of Schiff base complexes in catalysis (Cozzi, 2004). M.M. Abd-Elzaher reported the synthesis and anticancer activity of Schiff base complex with thiazole moiety (Abd-Elzaher et al., 2016). P. Kavitha et al studied the biological activity and DNA cleavage of Co (II) Schiff's base complexes (Kavitha et al., 2016).

In our pursuit to explore the field of bioinorganic chemistry we have synthesized thiazolyl Schiff bases by condensation of 2-(4-chlorophenyl)-4-methylthiazole-5-carbohydrazide and substituted aromatic aldehydes. Their Cu(II) and Ni(II) complexes were prepared and characterized by sophisticated spectroscopic techniques such as UV, FTIR, <sup>1</sup>H NMR and XRD.

## EXPERIMENTAL:

The chemicals used were of AR grade. Melting Points were determined by open capillary method and are uncorrected. IR spectra were recorded on a FT-IR spectrophotometer RZX (Perkin Elmer) and Mass spectra were recorded on a Q-TOF MICRO WATER, MS ES+3.79e3. <sup>1</sup>H spectra on a BRUKER AVANCE NEO 500 NMR spectrometer with DMSO-*d*<sub>6</sub> as a solvent and chemical shift (δ) are expressed in ppm using TMS as internal standard. UV-Visible spectra were recorded on Shimadzu UV-1800 instrument.

### *Synthesis of thiazolyl Schiff base ligand*

Equimolar mixture of 2-(4-chlorophenyl)-4-methylthiazole-5-carbohydrazide (0.01 mole) **3** and aromatic aldehyde **4** (0.01 mole) with 3 drops of Conc.H<sub>2</sub>SO<sub>4</sub> were refluxed in alcohol about 1-2 hrs (checked by TLC). After cooling, solid product thus obtained, it was filtered and recrystallized in alcohol to get Schiff base **5**. [Figure 1]

### **5D: (E)-N'-(2-chlorobenzylidene)-2-(4-chlorophenyl)-4-methylthiazole-5-carbohydrazide.**

IR (KBr): 3245 (N-H), 3058 (C-H), 1653 (C=O), 1090 (Ar-Cl), 832 (disubstituted benzene) cm<sup>-1</sup>; Mass: *m/z* 390 (M+1)<sup>+</sup>; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>): δ 2.77 (s, 3H, CH<sub>3</sub>), 7.48 (d, 2H, Ar-H), 7.54 (dd, 1H, Ar-H), 7.62 (d, 2H, Ar-H),

8.03 (m, 3H, Ar-H), 8.55 (s, 1H, N=C-H), 12.08 (s, 1H, NH); Elemental Analysis, Calculated: C<sub>18</sub>H<sub>13</sub>OCl<sub>2</sub>N<sub>3</sub>S: C, 55.39; H, 3.36; N, 10.77. Found: C, 55.37; H, 3.34; N, 10.75 %.

**5H: (E)-N<sup>o</sup>-(2-nitrobenzylidene)-2-(4-chlorophenyl)-4-methylthiazole-5-carbohydrazide.**

IR (KBr): 3165 (N-H), 3051(C-H), 1657 (C=O), 1528 (Ar-NO<sub>2</sub>), 1092 (Ar-Cl), 832 (disubstituted benzene) cm<sup>-1</sup>; Mass: *m/z* 401.29 (M+1)<sup>+</sup>; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>): δ 2.76 (s, 3H, CH<sub>3</sub>), 7.60 (d, 2H, Ar-H), 7.70 (dd, 1H, Ar-H), 7.89 (s, 1H, N=C-H), 8.02 (d, 1H, Ar-H) 8.11 (dd, 1H, Ar-H), 8.14 (d, 1H, Ar-H), 12.18 (s, 1H, NH); Elemental Analysis, Calculated: C<sub>18</sub>H<sub>13</sub>O<sub>3</sub>ClN<sub>4</sub>S: C, 53.94; H, 3.27; N, 13.98. Found: C, 53.97; H, 3.29; N, 13.95 %.

**Synthesis of Schiff base metal (II) complex**

A solution of metal salt (0.01 mole) with 15 ml of ethyl alcohol is added to a solution of (0.01 mole) Schiff base in 10 ml of ethyl alcohol with stirring. The reaction mixture was heated under reflux for 1-1.5 hr. The solid thus obtained was filtered off and then dried, to get Schiff base metal complex. [Figure 2]

**RESULTS AND DISCUSSION:**

The Schiff base ligand (5) was prepared from the condensation of 2-(4-chlorophenyl)-4-methylthiazole-5-carbohydrazide and substituted aromatic aldehydes in good yield [Table 1]. The ligand and its metal complexes are stable at room temperature.

FTIR analysis showed a strong band a 1657 cm<sup>-1</sup> was assigned to amide carbonyl group. The band at 1528 cm<sup>-1</sup> was attributed to nitro group and bands at 3165 cm<sup>-1</sup> and 3051 cm<sup>-1</sup> are allocated to the NH and CH stretching.

<sup>1</sup>HNMR shown singlet at 2.77 ppm for methyl group, peaks at 7.48, 7.54, 7.62 and 8.03 are for aromatic protons and most down field proton at 8.55 ppm accounts for azomethine (N=C-H) proton. The broad singlet at 12.08 ppm is due to NH group.

Mass spectra of ligand shown the molecular ion peak at 429.1 [M+1]

UV absorption of Cu (II) and Ni (II) metal complexes shown at the bands λ<sub>max</sub> at 325 and 328 nm respectively. The bands are assigned to n-π\* and charge transfer of azomethine group and thiazole ring nitrogen atom. The magnetic moment of Cu and Ni complex are 1.73 B.M. and 2.82 B.M. indicates octahedral geometries for both complexes (Gull & Hashmi, 2017).

Powder X-ray analysis of Cu (II) and Ni (II) metal complexes shown that the complexes are crystalline in nature. X-Ray diffractogram shows 25 reflections with maxima at 25.58 Å and 26.17 Å for Cu (II) and Ni (II) metal complexes respectively (Gull & Hashmi, 2017). [Figure 3 and 4]

**CONCLUSION:**

We have synthesized Schiff base ligand in good yield. Spectroscopic techniques FTIR, mass and <sup>1</sup>HNMR confirms its formation whereas UV and XRD reveals that the Cu (II) and Ni (II) metal complexes are crystalline in nature and geometries are octahedral for both the complexes. In conclusion, the versatile Cu (II) and Ni (II) metal complexes have excellent properties and worth for further studies on its biological activities.

**ACKNOWLEDGEMENT:**

Authors thanks to the Principal and Head, Department of Chemistry, Tuljaram Chaturchand College Baramati, Radhabai Kale Mahila Mahavidyalaya, Ahmednagar and New Arts, Commerce and Science College, Parner, Ahmednagar for providing necessary laboratory facilities and chemicals. Authors also thank to the Director, SAIF, Punjab University Chandigarh for FTIR, <sup>1</sup>HNMR and XRD analysis.

**REFERENCES:**

1. Abd-Elzaher M. M., Labib A. A., Mousa, H. A., Moustafa S. A., Ali, M. M. & El-Rashedy A. A., (2016) Synthesis, anticancer activity and molecular docking study of Schiff base complexes containing thiazole moiety, *Beni-Suef University Journal of Basic and Applied Sciences*, 5(1), 85-96. <https://doi.org/https://doi.org/10.1016/j.bjbas.2016.01.001>
2. Alsharif Z. A. & Alam M. A., (2017) Modular synthesis of thiazoline and thiazole derivatives by using a cascade protocol [10.1039/C7RA05993K], *RSC Advances*, 7(52), 32647-32651. <https://doi.org/10.1039/C7RA05993K>
3. Anacona J. R., Santaella J., Al-Shemary R. K. R., Amenta, J., Otero A., Ramos C. & Celis F., (2021) Ceftriaxone-based Schiff base transition metal(II) complexes, synthesis, characterization, bacterial toxicity, and DFT calculations. Enhanced antibacterial activity of a novel Zn(II) complex against *S. aureus* and *E. coli*. *J Inorg Biochem*, 223, 111519. <https://doi.org/10.1016/j.jinorgbio.2021.111519>
4. Bursal E., Turkan F., Buldurun K., Turan N., Aras A., Colak N., Murahari M. & Yergeri M. C., (2021) Transition metal complexes of a multidentate Schiff base ligand containing pyridine: synthesis, characterization, enzyme inhibitions, antioxidant properties and molecular docking studies, *Biometals*, 34(2), 393-406. <https://doi.org/10.1007/s10534-021-00287-z>
5. Chaviara A. T., Cox P. J., Repana K. H., Papi R. M., Papazisis K. T., Zambouli D., Kortsaris A. H., Kyriakidis D. A. & Bolos C. A., (2004) Copper(II) Schiff base coordination compounds of dien with heterocyclic aldehydes and 2-amino-5-methyl-thiazole: synthesis, characterization, antiproliferative and antibacterial studies. Crystal structure of CudienOOC12, *J Inorg Biochem*, 98(8), 1271-1283. <https://doi.org/10.1016/j.jinorgbio.2004.05.010>
6. Cozzi P. G., (2004) Metal–Salen Schiff base complexes in catalysis: practical aspects [10.1039/B307853C], *Chemical Society Reviews*, 33(7), 410-421. <https://doi.org/10.1039/B307853C>
7. Damercheli M., Mahdi M., Mehravi B. & Shafiee Ardestani M., (2019) Bioactive Salen-type Schiff Base Transition Metal Complexes as Possible Anticancer Agents, *Iran J Pharm Res*, 18(4), 2055-2066. <https://doi.org/10.22037/ijpr.2019.12792.11151>
8. Gull P. & Hashmi A. A., (2017) Synthesis, XRD and spectroscopic characterization of pharmacologically active Cu(II) and Zn(II) complexes, *Journal of Molecular Structure*, 1139, 264-268. <https://doi.org/https://doi.org/10.1016/j.molstruc.2017.03.053>
9. Karatepe M. & Karatas F., (2006) Antioxidant, pro-oxidant effect of the thiosemicarbazone derivative Schiff base (4-(1-phenylmethylcyclobutane-3-yl)-2-(2-hydroxybenzylidenehydrazino) thiazole) and its metal complexes on rats, *Cell Biochem Funct*, 24(6), 547-554. <https://doi.org/10.1002/cbf.1266>
10. Kavitha P., Rama Chary M., Singavarapu B. V. V. A. & Laxma Reddy K., (2016) Synthesis, characterization, biological activity and DNA cleavage studies of tridentate Schiff bases and their Co(II) complexes, *Journal of Saudi Chemical Society*, 20(1), 69-80. <https://doi.org/https://doi.org/10.1016/j.jscs.2013.03.005>
11. Manimohan M., Paulpandiyan R., Pugalmani S. & Sithique M. A., (2020) Biologically active Co (II), Cu (II), Zn (II) centered water soluble novel isoniazid grafted O-carboxymethyl chitosan Schiff base ligand metal

- complexes: synthesis, spectral characterisation and DNA nuclease activity, *Int J Biol Macromol*, 163, 801-816. <https://doi.org/10.1016/j.ijbiomac.2020.06.278>
12. Pontiki E., Hadjipavlou-Litina D. & Chaviara A. T., (2008) Evaluation of anti-inflammatory and antioxidant activities of copper (II) Schiff mono-base and copper(II) Schiff base coordination compounds of dien with heterocyclic aldehydes and 2-amino-5-methyl-thiazole, *J Enzyme Inhib Med Chem*, 23(6), 1011-1017. <https://doi.org/10.1080/14756360701841251>
13. Sahin O., Ozdemir U. O., Seferoglu N., Genc Z. K., Kaya K., Aydiner B., Tekin S. & Seferoglu Z., (2018) New platinum (II) and palladium (II) complexes of coumarin-thiazole Schiff base with a fluorescent chemosensor properties: Synthesis, spectroscopic characterization, X-ray structure determination, in vitro anticancer activity on various human carcinoma cell lines and computational studies, *J Photochem Photobiol B*, 178, 428-439. <https://doi.org/10.1016/j.jphotobiol.2017.11.030>
14. Shi S., Yu S., Quan L., Mansoor M., Chen Z., Hu H., Liu D., Liang Y. & Liang, F., (2020) Synthesis and antitumor activities of transition metal complexes of a bis-Schiff base of 2-hydroxy-1-naphthalenecarboxaldehyde, *J Inorg Biochem*, 210, 111173. <https://doi.org/10.1016/j.jinorgbio.2020.111173>
15. Singh A. & Barman P., (2021) Recent Advances in Schiff Base Ruthenium Metal Complexes: Synthesis and Applications, *Top Curr Chem (Cham)*, 379(4), 29. <https://doi.org/10.1007/s41061-021-00342-w>
16. Yernale N. G. & Bennikallu Hire Mathada M., (2014) Synthesis, characterization, antimicrobial, DNA cleavage, and in vitro cytotoxic studies of some metal complexes of schiff base ligand derived from thiazole and quinoline moiety, *Bioinorg Chem Appl*, 2014, 314963. <https://doi.org/10.1155/2014/314963>




Figure 1: Synthesis of Schiff base ligand




Figure 2: Synthesis of Schiff base metal complexes




Figure 3: X-Ray diffractogram of Cu (II) complex




Figure 4: X-Ray diffractogram of Ni (II) complex

Table-

1:

Comps	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	M.P. (°C)	Time (min.)	Yield (%)
<b>5A</b>	H	H	Cl	278-280	80	66
<b>5B</b>	H	H	Br	286-288	80	63
<b>5C</b>	H	H	OH	266-268	90	68
<b>5D</b>	Cl	H	H	280-282	80	65
<b>5E</b>	H	H	H	250-252	85	70
<b>5F</b>	H	H	OCH <sub>3</sub>	260-262	90	61
<b>5G</b>	H	H	NO <sub>2</sub>	288-290	75	77
<b>5H</b>	NO <sub>2</sub>	H	H	282-284	75	75

Physical data of synthesized Schiff bases